## Rotary embeddings of Praeger-Xu graphs

Luyi Liu

International center for mathematics, Sustech

September 17, 2025

### Мар

A map  $\mathcal{M}=(V,E,F)$  is a 2-cell embedding of a graph  $\Gamma=(V,E)$  into a closed surface  $\mathcal{S}.$ 

A map  $\mathcal{M}=(V,E,F)$  is a 2-cell embedding of a graph  $\Gamma=(V,E)$  into a closed surface  $\mathcal{S}.$ 




A map  $\mathcal{M} = (V, E, F)$  is a 2-cell embedding of a graph  $\Gamma = (V, E)$  into a closed surface  $\mathcal{S}$ .



#### The face set:

The open disks  $F := S \setminus \Gamma$ .

A map  $\mathcal{M} = (V, E, F)$  is a 2-cell embedding of a graph  $\Gamma = (V, E)$  into a closed surface  $\mathcal{S}$ .



#### The face set:


The open disks  $F := S \setminus \Gamma$ .

#### The flag set:


 $\Phi := \{(\alpha, e, f) : \alpha \in V, e \in E, f \in F \text{ are mutually incident}\}.$ 

Remark:  $|\Phi| = 4|E|$ .

## Automorphisms of maps



## Automorphisms of maps



#### Automorphisms of map

 $Aut\mathcal{M} := \{ \sigma \in Sym(\Phi) : \sigma \text{ preserves incidences between flags} \}.$ 

- **②**  $\operatorname{Aut}(\mathcal{M})_{\omega}$  is cyclic or dihedral,  $\omega \in V \cup E \cup F$ .  $\Rightarrow G_e \leqslant D_4$ .



 $(\alpha, e, f)$ : a flag.

## Rotary maps: a class of arc-regular map

 $(\alpha, e, f)$ : a flag.

## G-arc-transitive maps:

 $G \leqslant \operatorname{Aut}(\mathcal{M})$  is transitive on arcs of  $\mathcal{M}$ .

 $(\alpha, e, f)$ : a flag.

#### G-arc-transitive maps:

 $G \leqslant \operatorname{Aut}(\mathcal{M})$  is transitive on arcs of  $\mathcal{M}$ .

## G-vertex (face)-rotary:

 $G_{\alpha}$  ( $G_{f}$ ) induces a transitive cyclic subgroup on edges that incident with the vertex  $\alpha$  (the face f).

#### G-rotary map:

A map which is both *G*-vertex-rotary and *G*-face-rotary.

### Constructions of rotary maps

G: a finite group.

<sup>&</sup>lt;sup>0</sup>[15] C. H. Li, C. E. Praeger, and S. J. Song. Locally finite vertex-rotary maps and coset graphs with finite valency and finite edge multiplicity. *Journal of Combinatorial Theory. Series B*, 2024.

G: a finite group.

## $\mathsf{RotaMap}(\mathsf{G}, \rho, \tau)$

A rotary pair  $(\rho, \tau) \in G \times G$  of a group G satisfies  $G = \langle \rho, \tau \rangle$  and  $|\tau| = 2$ . Define an incidence configuration RotaMap $(G, \rho, \tau)$  by

vertex set  $[G:\langle \rho \rangle]$ , edge set  $[G:\langle \tau \rangle]$  and face set  $[G:\langle \rho \tau \rangle]$ ,

where two objects are incident if and only if their set intersection is non-empty.

<sup>&</sup>lt;sup>0</sup>[15] C. H. Li, C. E. Praeger, and S. J. Song. Locally finite vertex-rotary maps and coset graphs with finite valency and finite edge multiplicity. *Journal of Combinatorial Theory. Series B*, 2024.

G: a finite group.

## $\mathsf{RotaMap}(\mathsf{G}, \rho, \tau)$

A rotary pair  $(\rho, \tau) \in G \times G$  of a group G satisfies  $G = \langle \rho, \tau \rangle$  and  $|\tau| = 2$ . Define an incidence configuration RotaMap $(G, \rho, \tau)$  by

vertex set  $[G:\langle \rho \rangle]$ , edge set  $[G:\langle \tau \rangle]$  and face set  $[G:\langle \rho \tau \rangle]$ ,

where two objects are incident if and only if their set intersection is non-empty.

## [15, Proposition 5.1]

A G-rotary map is isomorphic to RotaMap $(G, \rho, \tau)$  for some rotary pair  $(\rho, \tau)$  for G.

### [15, Proposition 4.1]

Two maps RotaMap( $G, \rho, \tau$ ) and RotaMap( $H, \rho', \tau'$ ) are isomorphic if there is a group isomorphism  $f: G \to H$  such that  $f(\rho) = \rho'$  and  $f(\tau) = \tau'$ .

<sup>&</sup>lt;sup>0</sup>[15] C. H. Li, C. E. Praeger, and S. J. Song. Locally finite vertex-rotary maps and coset graphs with finite valency and finite edge multiplicity. *Journal of Combinatorial Theory. Series B*, 2024.

- $\mathcal{M} = \mathsf{RotaMap}(\mathsf{G}, \rho, \tau)$ .
- $M \lhd G$  with  $\rho, \tau \notin M$

 $<sup>^{0}</sup>$ [1] J. Chen, W. Fan, C. H. Li, and Y. Z. Zhu. Coverings of groups, regular dessins, and surfaces, 2024.

- $\mathcal{M} = \mathsf{RotaMap}(G, \rho, \tau)$ .
- $M \triangleleft G$  with  $\rho, \tau \notin M$

## Quotient rotary maps[1]

 $\mathcal{M}/M := \mathsf{RotaMap}(G/M, \rho M, \tau M)$ 

<sup>&</sup>lt;sup>0</sup>[1] J. Chen, W. Fan, C. H. Li, and Y. Z. Zhu. Coverings of groups, regular dessins, and surfaces, 2024.

- $\mathcal{M} = \mathsf{RotaMap}(G, \rho, \tau)$ .
- $M \triangleleft G$  with  $\rho, \tau \notin M$

## Quotient rotary maps[1]

 $\mathcal{M}/M := \mathsf{RotaMap}(G/M, \rho M, \tau M)$ 

- $\mathcal{M}_i = \mathsf{RotaMap}(G_i, \rho_i, \tau_i) \ (i = 1, \dots, n).$
- $H = \langle (\rho_1, \ldots, \rho_n), (\tau_1, \ldots, \tau_n) \rangle \leqslant \prod_{i=1}^n G_i$ .

<sup>&</sup>lt;sup>0</sup>[1] J. Chen, W. Fan, C. H. Li, and Y. Z. Zhu. Coverings of groups, regular dessins, and surfaces, 2024.

- $\mathcal{M} = \mathsf{RotaMap}(G, \rho, \tau)$ .
- $M \triangleleft G$  with  $\rho, \tau \notin M$

#### Quotient rotary maps[1]

 $\mathcal{M}/M := \mathsf{RotaMap}(G/M, \rho M, \tau M)$ 

- $\mathcal{M}_i = \mathsf{RotaMap}(G_i, \rho_i, \tau_i) \ (i = 1, \dots, n).$
- $H = \langle (\rho_1, \ldots, \rho_n), (\tau_1, \ldots, \tau_n) \rangle \leqslant \prod_{i=1}^n G_i$ .

### Direct products

$$\prod_{i=1}^n \mathcal{M}_i := \mathsf{RotaMap}(H, (\rho_1, \dots, \rho_n), (\tau_1, \dots, \tau_n)).$$

 $<sup>^{0}</sup>$ [1] J. Chen, W. Fan, C. H. Li, and Y. Z. Zhu. Coverings of groups, regular dessins, and surfaces, 2024.

#### Praeger-Xu graphs

## Praeger-Xu graph


Let p, r, s be positive integers such that  $p\geqslant 2$  and  $r\geqslant 3$ . Define a simple graph C(p,r,s)=(V,E) as follows:

- ① the vertex set V is  $\mathbb{Z}_r \times \mathbb{Z}_p^s$ ;
- $\bullet$  the edge set E is defined to be the set of all pairs of the form

$$\{(i, x_0, x_1, \ldots, x_{s-1}), (i+1, x_1, \ldots, x_{s-1}, x_s)\}$$

for every  $i \in \mathbb{Z}_r$  and  $x_0, x_1, \ldots, x_{s-1}, x_s \in \mathbb{Z}_p$ .

There are  $p^{s+1}r$  edges.



#### Praeger-Xu graphs

## Praeger-Xu graph

Let p, r, s be positive integers such that  $p \geqslant 2$  and  $r \geqslant 3$ . Define a simple graph C(p, r, s) = (V, E) as follows:

- ① the vertex set V is  $\mathbb{Z}_r \times \mathbb{Z}_p^s$ ;
- $\bullet$  the edge set E is defined to be the set of all pairs of the form

$$\{(i, x_0, x_1, \ldots, x_{s-1}), (i+1, x_1, \ldots, x_{s-1}, x_s)\}$$

for every  $i \in \mathbb{Z}_r$  and  $x_0, x_1, \ldots, x_{s-1}, x_s \in \mathbb{Z}_p$ .

There are  $p^{s+1}r$  edges.



#### Augmented PX-graphs

PX-graphs, multicycles  $C_r^{(p)}$  (denoted by  $C^*(p, r, 0, 1)$ ), and cycles  $C_{pr}$  (denoted by  $C^*(p, r, 0, -1)$ ).

### Properties of Praeger-Xu graphs

## [19, Theorem 2.10]

The graph C(p, r, s) is symmetric if and only if  $r \ge s + 1$ , and is vertex transitive if and only if  $r \ge s$ .

<sup>&</sup>lt;sup>0</sup>19. C. E. Praeger and M. Y. Xu. A Characterization of a Class of Symmetric Graphs of Twice Prime Valency. European Journal of Combinatorics, 1989

## Properties of Praeger-Xu graphs

#### [19, Theorem 2.10]

The graph C(p, r, s) is symmetric if and only if  $r \ge s + 1$ , and is vertex transitive if and only if  $r \ge s$ .

#### [19, Theorem 2.13]

The full automorphism group of C(p, r, s) is equal to  $S_p$  wr  $D_{2r}$  where  $(r, s) \neq (4, 1)$ ,  $r \ge \max\{s + 1, 3\}$  and p is odd.

<sup>&</sup>lt;sup>0</sup>19. C. E. Praeger and M. Y. Xu. A Characterization of a Class of Symmetric Graphs of Twice Prime Valency. European Journal of Combinatorics, 1989

### Properties of Praeger-Xu graphs

#### [19, Theorem 2.10]

The graph C(p, r, s) is symmetric if and only if  $r \ge s + 1$ , and is vertex transitive if and only if  $r \ge s$ .

### [19, Theorem 2.13]

The full automorphism group of C(p,r,s) is equal to  $S_p$  wr  $D_{2r}$  where  $(r,s) \neq (4,1)$ ,  $r \geqslant \max\{s+1,3\}$  and p is odd.

## [19, Theorem 1]

Let  $\Gamma$  be a connected, simple, G-arc-transitive graph of valency 2p. If G contains an abelian normal p-subgroup which is not semiregular on the vertices of  $\Gamma$ , then  $\Gamma = C(p,r,s)$  for some  $r \geqslant \max\{3,s+1\}$  and  $s \geqslant 1$ .

<sup>&</sup>lt;sup>0</sup>19. C. E. Praeger and M. Y. Xu. A Characterization of a Class of Symmetric Graphs of Twice Prime Valency. European Journal of Combinatorics, 1989

Let  $s \geqslant 0$ ,  $r \geqslant \max\{s+1,3\}$  and  $(r,s) \neq (4,1)$ .

Let  $s \ge 0$ ,  $r \ge \max\{s+1,3\}$  and  $(r,s) \ne (4,1)$ .

## [2, Lemma 4.5]

Suppose that G is an arc-regular subgroup of  $\operatorname{Aut}(\operatorname{C}(p,r,s))$ . Then

$$G\cong \mathbb{Z}_p^{s+1}$$
: $\mathrm{D}_{2r}$ .

Let  $s \ge 0$ ,  $r \ge \max\{s+1,3\}$  and  $(r,s) \ne (4,1)$ .

## [2, Lemma 4.5]

Suppose that G is an arc-regular subgroup of  $\operatorname{Aut}(\operatorname{C}(p,r,s))$ . Then

$$G\cong \mathbb{Z}_p^{s+1}$$
: $\mathrm{D}_{2r}$ .

#### Proof

The order  $|G| = 2p^{s+1}r$  which is the arc number of C(p, r, s).

Let  $s \ge 0$ ,  $r \ge \max\{s+1,3\}$  and  $(r,s) \ne (4,1)$ .

## [2, Lemma 4.5]

Suppose that G is an arc-regular subgroup of  $\operatorname{Aut}(\operatorname{C}(p,r,s))$ . Then

$$G\cong \mathbb{Z}_p^{s+1}$$
: $\mathrm{D}_{2r}$ .

#### Proof

The order  $|G| = 2p^{s+1}r$  which is the arc number of C(p, r, s). Let  $\overline{\Gamma} = C(p, r, s)_{S_p^r}$  which is isomorphic to  $C_r$ .

Let  $s \ge 0$ ,  $r \ge \max\{s+1,3\}$  and  $(r,s) \ne (4,1)$ .

#### [2, Lemma 4.5]

Suppose that G is an arc-regular subgroup of  $\operatorname{Aut}(\operatorname{C}(p,r,s))$ . Then

$$G\cong \mathbb{Z}_p^{s+1}$$
: $\mathrm{D}_{2r}$ .

#### Proof

The order  $|G|=2p^{s+1}r$  which is the arc number of  $\mathrm{C}(p,r,s)$ . Let  $\overline{\Gamma}=\mathrm{C}(p,r,s)_{\mathrm{S}_p^r}$  which is isomorphic to  $\mathrm{C}_r$ . Since G is transitive on arcs of  $\overline{\Gamma}\cong\mathrm{C}_r$  with the kernel  $G\cap\mathrm{S}_p^r$ , we have that

$$G/(G \cap S_p^r) \cong D_{2r} \text{ and } |G \cap S_p^r| = p^{s+1}.$$

Let  $s \ge 0$ ,  $r \ge \max\{s+1,3\}$  and  $(r,s) \ne (4,1)$ .

#### [2, Lemma 4.5]

Suppose that G is an arc-regular subgroup of  $\operatorname{Aut}(\operatorname{C}(p,r,s))$ . Then

$$G\cong \mathbb{Z}_p^{s+1}$$
:D<sub>2r</sub>.

#### Proof

The order  $|G|=2p^{s+1}r$  which is the arc number of  $\mathrm{C}(p,r,s)$ . Let  $\overline{\Gamma}=\mathrm{C}(p,r,s)_{\mathrm{S}_p^r}$  which is isomorphic to  $\mathrm{C}_r$ . Since G is transitive on arcs of  $\overline{\Gamma}\cong\mathrm{C}_r$  with the kernel  $G\cap\mathrm{S}_p^r$ , we have that

$$G/(G \cap S_p') \cong D_{2r}$$
 and  $|G \cap S_p'| = p^{s+1}$ .

It follows from  $(S_p^r)_p = \mathbb{Z}_p^r$  that

$$G=\mathbb{Z}_p^{s+1}$$
: $\mathrm{D}_{2r}$ .

 $\mathrm{C}(\textit{p},4,1)$  is the complete bipartite graph  $\mathrm{K}_{2\textit{p},2\textit{p}}.$ 

C(p, 4, 1) is the complete bipartite graph  $K_{2p,2p}$ .  $Aut(K_{2p,2p}) = S_{2p}$  wr  $\mathbb{Z}_2$ .

## [2, Lemma 4.6]

Suppose that G is an arc-regular subgroup of  $\operatorname{Aut}(\mathrm{K}_{2\rho,2\rho})$ . Then  $G\cong\mathbb{Z}_{\rho}^2:\mathrm{D}_8.$ 

#### Proof.

The order of G is  $8p^2$ .

C(p, 4, 1) is the complete bipartite graph  $K_{2p,2p}$ .  $Aut(K_{2p,2p}) = S_{2p}$  wr  $\mathbb{Z}_2$ .

#### [2, Lemma 4.6]

Suppose that G is an arc-regular subgroup of  $\operatorname{Aut}(K_{2p,2p})$ . Then  $G \cong \mathbb{Z}_p^2:D_8$ .

#### Proof.

The order of G is  $8p^2$ . Let L is the subgroup which preserving the two parts of  $K_{2p,2p}$ , and so  $|L|=4p^2$ . Note that  $L=G_{\alpha}G_{\beta}$ , where  $\alpha,\beta$  are two vertices in the two parts, respectively.

C(p,4,1) is the complete bipartite graph  $K_{2p,2p}$ .  $Aut(K_{2p,2p}) = S_{2p}$  wr  $\mathbb{Z}_2$ .

#### [2, Lemma 4.6]

Suppose that G is an arc-regular subgroup of  $\operatorname{Aut}(\mathrm{K}_{2p,2p})$ . Then  $G\cong\mathbb{Z}_p^2:\mathrm{D}_8$ .

#### Proof.

The order of G is  $8p^2$ . Let L is the subgroup which preserving the two parts of  $K_{2p,2p}$ , and so  $|L|=4p^2$ . Note that  $L=G_{\alpha}G_{\beta}$ , where  $\alpha,\beta$  are two vertices in the two parts, respectively. Since  $G_{\alpha}\cong G_{\beta}\cong D_{2p}$  or  $\mathbb{Z}_{2p}$ , we have that  $L_p\cong \mathbb{Z}_p^2$ .

C(p, 4, 1) is the complete bipartite graph  $K_{2p,2p}$ .  $Aut(K_{2p,2p}) = S_{2p}$  wr  $\mathbb{Z}_2$ .

### [2, Lemma 4.6]

Suppose that G is an arc-regular subgroup of  $\operatorname{Aut}(\mathrm{K}_{2\rho,2\rho})$ . Then  $G\cong\mathbb{Z}_p^2:\mathrm{D}_8$ .

#### Proof.

The order of G is  $8p^2$ . Let L is the subgroup which preserving the two parts of  $K_{2p,2p}$ , and so  $|L|=4p^2$ . Note that  $L=G_{\alpha}G_{\beta}$ , where  $\alpha,\beta$  are two vertices in the two parts, respectively. Since  $G_{\alpha}\cong G_{\beta}\cong D_{2p}$  or  $\mathbb{Z}_{2p}$ , we have that  $L_p\cong \mathbb{Z}_p^2$ .

We claim that  $L_p \lhd L$ , and so  $L \cong \mathbb{Z}_p^2: \mathbb{Z}_2^2$  implying

$$G \cong (\mathbb{Z}_p^2:\mathbb{Z}_2^2):\mathbb{Z}_2 \cong \mathbb{Z}_p^2:D_8.$$

C(p, 4, 1) is the complete bipartite graph  $K_{2p,2p}$ .  $Aut(K_{2p,2p}) = S_{2p} \ wr \ \mathbb{Z}_2$ .

#### [2, Lemma 4.6]

Suppose that G is an arc-regular subgroup of  $\operatorname{Aut}(\mathrm{K}_{2p,2p})$ . Then  $G\cong\mathbb{Z}_p^2:\mathrm{D}_8$ .

#### Proof.

The order of G is  $8p^2$ . Let L is the subgroup which preserving the two parts of  $K_{2p,2p}$ , and so  $|L|=4p^2$ . Note that  $L=G_{\alpha}G_{\beta}$ , where  $\alpha,\beta$  are two vertices in the two parts, respectively. Since  $G_{\alpha}\cong G_{\beta}\cong D_{2p}$  or  $\mathbb{Z}_{2p}$ , we have that  $L_p\cong \mathbb{Z}_p^2$ .

We claim that  $L_p \lhd L$ , and so  $L \cong \mathbb{Z}_p^2 : \mathbb{Z}_2^2$  implying

$$G\cong (\mathbb{Z}_p^2{:}\mathbb{Z}_2^2){:}\mathbb{Z}_2\cong \mathbb{Z}_p^2{:}\mathrm{D}_8.$$

Let p=3. By Brodkey's theorem, there are two Sylow 3-subgroups P and Q of L such that  $P\cap Q=\mathrm{O}_3(L)\cong \mathbb{Z}_3$ . The quotient group  $L/\mathrm{O}_3(L)\cong \mathrm{A}_4$  since its Sylow 3-subgroups are not normal.

C(p, 4, 1) is the complete bipartite graph  $K_{2p,2p}$ .  $Aut(K_{2p,2p}) = S_{2p}$  wr  $\mathbb{Z}_2$ .

#### [2, Lemma 4.6]

Suppose that G is an arc-regular subgroup of  $\operatorname{Aut}(\mathrm{K}_{2p,2p})$ . Then  $G\cong\mathbb{Z}_p^2:\mathrm{D}_8$ .

#### Proof.

The order of G is  $8p^2$ . Let L is the subgroup which preserving the two parts of  $K_{2p,2p}$ , and so  $|L|=4p^2$ . Note that  $L=G_{\alpha}G_{\beta}$ , where  $\alpha,\beta$  are two vertices in the two parts, respectively. Since  $G_{\alpha}\cong G_{\beta}\cong D_{2p}$  or  $\mathbb{Z}_{2p}$ , we have that  $L_p\cong \mathbb{Z}_p^2$ .

We claim that  $L_p \triangleleft L$ , and so  $L \cong \mathbb{Z}_p^2 : \mathbb{Z}_2^2$  implying

$$G\cong (\mathbb{Z}_p^2{:}\mathbb{Z}_2^2){:}\mathbb{Z}_2\cong \mathbb{Z}_p^2{:}\mathrm{D}_8.$$

Let p=3. By Brodkey's theorem, there are two Sylow 3-subgroups P and Q of L such that  $P\cap Q=\mathrm{O}_3(L)\cong\mathbb{Z}_3$ . The quotient group  $L/\mathrm{O}_3(L)\cong\mathrm{A}_4$  since its Sylow 3-subgroups are not normal. Note that  $L/\mathrm{O}_3(L)=(G_\alpha G_\beta)/\mathrm{O}_3(L)$ . Contradiction comes to that  $G_\alpha\cong G_\beta\cong\mathbb{Z}_{2p}$  or  $\mathrm{D}_{2p}$ .

$$\operatorname{Aut}(C_{pr} = C^*(p, r, 0, -1)) = D_{2pr}, \operatorname{Aut}(C_r^{(p)} = C^*(p, r, 0, 1)) = \mathbb{Z}_p:D_{2r}.$$

$$\operatorname{Aut}(\mathrm{C}_{pr}=\mathrm{C}^*(p,r,0,-1))=\mathrm{D}_{2pr},\ \operatorname{Aut}(\mathrm{C}_r^{(p)}=\mathrm{C}^*(p,r,0,1))=\mathbb{Z}_p:\mathrm{D}_{2r}.$$

#### Theorem

Let  $s\geqslant 0$  be an integer, let  $\delta\in\{1,-1\}$  and let r be an integer such that  $r\geqslant \max\{3,s+1\}$ . Let G be an arc-regular group of automorphisms of  $C^*(p,r,s,\delta)$ . Then  $G\cong\mathbb{Z}_p^{s+1}\colon D_{2r}$ .

$$\operatorname{Aut}(C_{pr} = \operatorname{C}^*(p, r, 0, -1)) = \operatorname{D}_{2pr}, \ \operatorname{Aut}(\operatorname{C}_r^{(p)} = \operatorname{C}^*(p, r, 0, 1)) = \mathbb{Z}_p: \operatorname{D}_{2r}.$$

#### Theorem

Let  $s\geqslant 0$  be an integer, let  $\delta\in\{1,-1\}$  and let r be an integer such that  $r\geqslant \max\{3,s+1\}$ . Let G be an arc-regular group of automorphisms of  $C^*(p,r,s,\delta)$ . Then  $G\cong \mathbb{Z}_p^{s+1}\colon \mathsf{D}_{2r}$ .

# Corollary

Let  $\mathcal{M}$  be a G-rotary map with underlying graph isomorphic to  $C^*(p, r, s, \delta)$ . Then G is isomorphic to  $\mathbb{Z}_p^{s+1}: \mathsf{D}_{2r}$ .

$$\operatorname{Aut}(C_{\rho r} = C^*(\rho, r, 0, -1)) = D_{2\rho r}, \operatorname{Aut}(C_r^{(\rho)} = C^*(\rho, r, 0, 1)) = \mathbb{Z}_{\rho}: D_{2r}.$$

#### **Theorem**

Let  $s\geqslant 0$  be an integer, let  $\delta\in\{1,-1\}$  and let r be an integer such that  $r\geqslant \max\{3,s+1\}$ . Let G be an arc-regular group of automorphisms of  $C^*(p,r,s,\delta)$ . Then  $G\cong \mathbb{Z}_p^{s+1}\colon \mathsf{D}_{2r}$ .

# Corollary

Let  $\mathcal{M}$  be a G-rotary map with underlying graph isomorphic to  $C^*(p, r, s, \delta)$ . Then G is isomorphic to  $\mathbb{Z}_p^{s+1}$ :  $D_{2r}$ .

#### Lemma

Let  $G = \mathbb{Z}_p^{s+1}$ :  $D_{2r}$  with  $s \geqslant 0$ , and let  $\mathcal{M}$  be a G-arc-regular map. If  $|G_{\alpha}| = 2p$ , then the underlying graph  $\Gamma$  is isomorphic to  $C^*(p, r, s)$ .

$$\operatorname{Aut}(C_{pr} = \operatorname{C}^*(p, r, 0, -1)) = \operatorname{D}_{2pr}, \ \operatorname{Aut}(\operatorname{C}_r^{(p)} = \operatorname{C}^*(p, r, 0, 1)) = \mathbb{Z}_p: \operatorname{D}_{2r}.$$

#### **Theorem**

Let  $s\geqslant 0$  be an integer, let  $\delta\in\{1,-1\}$  and let r be an integer such that  $r\geqslant \max\{3,s+1\}$ . Let G be an arc-regular group of automorphisms of  $C^*(p,r,s,\delta)$ . Then  $G\cong \mathbb{Z}_p^{s+1}\colon D_{2r}$ .

## Corollary

Let  $\mathcal{M}$  be a G-rotary map with underlying graph isomorphic to  $C^*(p, r, s, \delta)$ . Then G is isomorphic to  $\mathbb{Z}_p^{s+1}$ :  $D_{2r}$ .

### Lemma

Let  $G = \mathbb{Z}_p^{s+1}$ :  $D_{2r}$  with  $s \geqslant 0$ , and let  $\mathcal{M}$  be a G-arc-regular map. If  $|G_{\alpha}| = 2p$ , then the underlying graph  $\Gamma$  is isomorphic to  $C^*(p, r, s)$ .

$$|G_{\alpha\beta}| = egin{cases} 2 \Rightarrow r = 2; \ p \Rightarrow ext{the underlying graph is the multi-cycle C}^*(p,r,0,1) \ 2p \Rightarrow r = 2; \end{cases}$$

A PX map is a map with the underlying graph is a augmented Praeger-Xu graph  $C^*(p, r, s, \delta)$ , the length of this map is defined as r.

A PX map is a map with the underlying graph is a augmented Praeger-Xu graph  $\mathrm{C}^*(p,r,s,\delta)$ , the length of this map is defined as r. Let  $\mathcal M$  be a G-rotary embedding of  $\mathrm{C}^*(p,r,s,\delta)$  with  $p\nmid r$ . By Lemma 11,  $G\cong \mathbb{Z}_{s}^{s+1}:_{\imath}\mathrm{D}_{2r}$ .

A PX map is a map with the underlying graph is a augmented Praeger-Xu graph  $\mathrm{C}^*(p,r,s,\delta)$ , the length of this map is defined as r. Let  $\mathcal M$  be a G-rotary embedding of  $\mathrm{C}^*(p,r,s,\delta)$  with  $p\nmid r$ . By Lemma 11,  $G\cong \mathbb{Z}_p^{s+1}:_{\psi}\mathrm{D}_{2r}$ .

## Irreducible G-rotary PX maps

The *G*-rotary PX map is *irreducible* if  $\psi$  is irreducible.

A PX map is a map with the underlying graph is a augmented Praeger-Xu graph  $\mathrm{C}^*(p,r,s,\delta)$ , the length of this map is defined as r. Let  $\mathcal M$  be a G-rotary embedding of  $\mathrm{C}^*(p,r,s,\delta)$  with  $p\nmid r$ . By Lemma 11,  $G\cong \mathbb Z_p^{s+1}:_\psi \mathrm{D}_{2r}$ .

### Irreducible G-rotary PX maps

The *G*-rotary PX map is *irreducible* if  $\psi$  is irreducible.

- $G = V:_{\psi} D = \mathbb{Z}_p^d:_{\psi} D_{2r}, d \geqslant 2, \psi$  is irreducible.
- $D_{2r} = \langle c \rangle : \langle b \rangle = \mathbb{Z}_r : \mathbb{Z}_2$ .
- $v^{b^{-1}} = \psi(b)(v)$ ,  $v \in V$  and  $b \in D$ .

Note that the degree d is even, and  $C_V(x) \cong \mathbb{Z}_p^{d/2}$  for each involution  $x \in D \setminus Z(D)[2$ , Lemma 2.9].

A PX map is a map with the underlying graph is a augmented Praeger-Xu graph  $\mathrm{C}^*(p,r,s,\delta)$ , the length of this map is defined as r. Let  $\mathcal M$  be a G-rotary embedding of  $\mathrm{C}^*(p,r,s,\delta)$  with  $p\nmid r$ . By Lemma 11,  $G\cong \mathbb Z_p^{s+1}:_\psi \mathrm{D}_{2r}$ .

### Irreducible G-rotary PX maps

The G-rotary PX map is irreducible if  $\psi$  is irreducible.

- $G = V:_{\psi}D = \mathbb{Z}_p^d:_{\psi}D_{2r}, d \geqslant 2, \psi$  is irreducible.
- $D_{2r} = \langle c \rangle : \langle b \rangle = \mathbb{Z}_r : \mathbb{Z}_2$ .
- $v^{b^{-1}} = \psi(b)(v)$ ,  $v \in V$  and  $b \in D$ .

Note that the degree d is even, and  $C_V(x) \cong \mathbb{Z}_p^{d/2}$  for each involution  $x \in D \setminus Z(D)[2$ , Lemma 2.9].

### Proposition

Let  $x, y \in D$  be involutions such that  $D = \langle x, y \rangle$ , and let  $v \neq v' \in C_V(x) \setminus \{1\}$ . Then RotaMap $(G, vx, y) \cong \text{RotaMap}(G, v'x, y)$  are rotary PX maps

Since  $\langle vx \rangle \cap V \cong \mathbb{Z}_p$ , the map RotaMap(G, vx, y) has underlying graph being a PX graph by Theorem 11.

Since  $\langle vx \rangle \cap V \cong \mathbb{Z}_p$ , the map RotaMap(G, vx, y) has underlying graph being a PX graph by Theorem 11. Note that there exists an  $\sigma \in (\operatorname{End}_D(V))^{\times}$  such that  $\sigma(v) = v'[2$ , Lemma 2.10].

Since  $\langle vx \rangle \cap V \cong \mathbb{Z}_p$ , the map RotaMap(G, vx, y) has underlying graph being a PX graph by Theorem 11. Note that there exists an  $\sigma \in (\operatorname{End}_D(V))^{\times}$  such that  $\sigma(v) = v'[2$ , Lemma 2.10]. Define  $f: G \to G$ ,  $f(wg) := \sigma(w)g$  for every  $w \in V$ ,  $g \in D$ .

Since  $\langle vx \rangle \cap V \cong \mathbb{Z}_p$ , the map RotaMap(G, vx, y) has underlying graph being a PX graph by Theorem 11. Note that there exists an  $\sigma \in (\operatorname{End}_D(V))^{\times}$  such that  $\sigma(v) = v'[2$ , Lemma 2.10]. Define  $f: G \to G$ ,  $f(wg) := \sigma(w)g$  for every  $w \in V$ ,  $g \in D$ . The function f is one-to-one,

Since  $\langle vx \rangle \cap V \cong \mathbb{Z}_p$ , the map RotaMap(G,vx,y) has underlying graph being a PX graph by Theorem 11. Note that there exists an  $\sigma \in (\operatorname{End}_D(V))^{\times}$  such that  $\sigma(v) = v'[2$ , Lemma 2.10]. Define  $f: G \to G$ ,  $f(wg) := \sigma(w)g$  for every  $w \in V$ ,  $g \in D$ . The function f is one-to-one, and is a group homomorphism as: for  $w_1g_1, w_2g_2 \in G$ ,

$$f(w_1g_1w_2g_2) = f(w_1w_2^{g_1^{-1}}g_1g_2)$$

$$= \sigma(w_1)\sigma(w_2^{g_1^{-1}})g_1g_2$$

$$= \sigma(w_1)(\sigma(w_2))^{g_1^{-1}}g_1g_2$$

$$= \sigma(w_1)g_1\sigma(w_2)g_2$$

$$= f(w_1g_1)f(w_2g_2).$$

By definition, RotaMap $(G, vx, y) \cong \text{RotaMap}(G, v'x, y)$ .

Since  $\langle vx \rangle \cap V \cong \mathbb{Z}_p$ , the map RotaMap(G, vx, y) has underlying graph being a PX graph by Theorem 11. Note that there exists an  $\sigma \in (\operatorname{End}_D(V))^{\times}$  such that  $\sigma(v) = v'[2$ , Lemma 2.10]. Define  $f: G \to G$ ,  $f(wg) := \sigma(w)g$  for every  $w \in V$ ,  $g \in D$ . The function f is one-to-one, and is a group homomorphism as: for  $w_1g_1, w_2g_2 \in G$ ,

$$f(w_1g_1w_2g_2) = f(w_1w_2^{g_1^{-1}}g_1g_2)$$

$$= \sigma(w_1)\sigma(w_2^{g_1^{-1}})g_1g_2$$

$$= \sigma(w_1)(\sigma(w_2))^{g_1^{-1}}g_1g_2$$

$$= \sigma(w_1)g_1\sigma(w_2)g_2$$

$$= f(w_1g_1)f(w_2g_2).$$

By definition, RotaMap(G, vx, y)  $\cong$  RotaMap(G, v'x, y).

#### $\rho_{x}$

For any  $x \in D \leq G = V_{:\psi}D$ , let  $\rho_x$  denote an element in the set

$$\{vx \mid 1 \neq v \in C_V(x)\}.$$

$$G = V:_{\psi}D = \mathbb{Z}_p^d:_{\psi}D_{2r}, \ d \geqslant 2, \ \psi \text{ is irreducible.}$$

$$G = V:_{\psi}D = \mathbb{Z}_{p}^{d}:_{\psi}D_{2r}, d \geqslant 2, \psi$$
 is irreducible.

### Proposition

Let  $x_1, y_1, x_2, y_2 \in D$  be involutions where  $D = \langle x_1, y_1 \rangle = \langle x_2, y_2 \rangle$ . Let  $\sigma \in \operatorname{Aut}(D)$  be given by  $\sigma(x_1) = x_2$  and  $\sigma(y_1) = y_2$ . Then

- $\blacksquare \ \, \mathsf{RotaMap}(\mathsf{G},\rho_{\mathsf{x}_1},y_1) \cong \mathsf{RotaMap}(\mathsf{G},\rho_{\mathsf{x}_2},y_2) \ \mathsf{if} \ \mathsf{and} \ \mathsf{only} \ \mathsf{if} \ \psi \cong \psi \circ \sigma.$
- **1** Define  $\mathcal{RM}_G := \{ \mathsf{RotaMap}(G, \rho_x, y) \mid \langle x, y \rangle = D \}$ . The mapping

$$\psi^{\operatorname{Aut}(D_{2r})} \to \mathcal{RM}_{\mathcal{G}},$$
$$\psi \circ \eta \mapsto \operatorname{RotaMap}(\mathcal{G}, \rho_{\eta(x_1)}, \eta(y_1))$$

is a one-to-one correspondence.

$$G = V:_{\psi} D = \mathbb{Z}_{p}^{d}:_{\psi} D_{2r}, d \geqslant 2, \psi$$
 is irreducible.

### Proposition

Let  $x_1, y_1, x_2, y_2 \in D$  be involutions where  $D = \langle x_1, y_1 \rangle = \langle x_2, y_2 \rangle$ . Let  $\sigma \in \operatorname{Aut}(D)$  be given by  $\sigma(x_1) = x_2$  and  $\sigma(y_1) = y_2$ . Then

- RotaMap $(G, \rho_{x_1}, y_1) \cong \text{RotaMap}(G, \rho_{x_2}, y_2)$  if and only if  $\psi \cong \psi \circ \sigma$ .
- ① Define  $\mathcal{RM}_G := \{ \mathsf{RotaMap}(G, \rho_x, y) \mid \langle x, y \rangle = D \}$ . The mapping

$$\psi^{\operatorname{Aut}(D_{2r})} \to \mathcal{RM}_{G},$$
$$\psi \circ \eta \mapsto \operatorname{RotaMap}(G, \rho_{\eta(x_{1})}, \eta(y_{1}))$$

is a one-to-one correspondence.

#### Proof

 $\Rightarrow$ : Let  $f \in \operatorname{Aut}(G)$  be the isomorphism. Since  $x_1$  and  $x_2$  are involutions, there is

$$f(x_1) = f(x_1^p) = f(\rho_{x_1}^p) = \rho_{x_2}^p = x_2.$$

So  $\sigma := f|_D \in \operatorname{Aut}(D)$ .

$$G = V:_{\psi} D = \mathbb{Z}_{p}^{d}:_{\psi} D_{2r}, d \geqslant 2, \psi$$
 is irreducible.

### Proposition

Let  $x_1, y_1, x_2, y_2 \in D$  be involutions where  $D = \langle x_1, y_1 \rangle = \langle x_2, y_2 \rangle$ . Let  $\sigma \in \operatorname{Aut}(D)$  be given by  $\sigma(x_1) = x_2$  and  $\sigma(y_1) = y_2$ . Then

- RotaMap $(G, \rho_{x_1}, y_1) \cong \text{RotaMap}(G, \rho_{x_2}, y_2)$  if and only if  $\psi \cong \psi \circ \sigma$ .
- **1** Define  $\mathcal{RM}_G := \{ \mathsf{RotaMap}(G, \rho_x, y) \mid \langle x, y \rangle = D \}$ . The mapping

$$\psi^{\operatorname{Aut}(D_{2r})} \to \mathcal{RM}_{\mathcal{G}},$$

$$\psi \circ \eta \mapsto \operatorname{\mathsf{RotaMap}}(\mathcal{G}, \rho_{\eta(\mathsf{x}_1)}, \eta(\mathsf{y}_1))$$

is a one-to-one correspondence.

#### Proof

 $\Rightarrow$ : Let  $f \in \operatorname{Aut}(G)$  be the isomorphism. Since  $x_1$  and  $x_2$  are involutions, there is

$$f(x_1) = f(x_1^p) = f(\rho_{x_1}^p) = \rho_{x_2}^p = x_2.$$

So  $\sigma := f|_D \in \operatorname{Aut}(D)$ . Straightforward checking shows that  $f|_V$  serves as the isomorphism between  $\psi \circ \sigma$  and  $\psi$ .

# Proof

 $\Leftarrow: \ \mathsf{Denote} \ \ell \ \mathsf{the} \ \mathsf{isomorphism} \ \mathsf{between} \ \psi \ \mathsf{and} \ \psi \circ \sigma.$ 

## Proof

 $\Leftarrow$ : Denote  $\ell$  the isomorphism between  $\psi$  and  $\psi \circ \sigma$ . Define  $f: G \to G$  by  $vg \mapsto \ell(v)\sigma(g)$ . Then  $f \in \operatorname{Aut}(G)$  is such that  $f(x_1) = x_2$  and  $f(y_1) = y_2$ . Statement i is proved.

#### Proof

 $\Leftarrow$ : Denote  $\ell$  the isomorphism between  $\psi$  and  $\psi \circ \sigma$ . Define  $f: G \to G$  by  $vg \mapsto \ell(v)\sigma(g)$ . Then  $f \in \operatorname{Aut}(G)$  is such that  $f(x_1) = x_2$  and  $f(y_1) = y_2$ . Statement i is proved.

Note that  $\operatorname{Aut}(D)$  acts transitively on the set  $\mathcal{RM}_G$ . There is  $|\mathcal{RM}_G| = |\operatorname{Aut}(D)|/|\operatorname{Aut}(D)_\psi| = |\psi^{\operatorname{Aut}(D)}|$ , which completes the proof.

#### Proof

 $\Leftarrow$ : Denote  $\ell$  the isomorphism between  $\psi$  and  $\psi \circ \sigma$ . Define  $f: G \to G$  by  $vg \mapsto \ell(v)\sigma(g)$ . Then  $f \in \operatorname{Aut}(G)$  is such that  $f(x_1) = x_2$  and  $f(y_1) = y_2$ . Statement i is proved.

Note that  $\operatorname{Aut}(D)$  acts transitively on the set  $\mathcal{RM}_G$ . There is  $|\mathcal{RM}_G| = |\operatorname{Aut}(D)|/|\operatorname{Aut}(D)_\psi| = |\psi^{\operatorname{Aut}(D)}|$ , which completes the proof.

We claim that the number of G-rotary augmented PX maps equals  $|\psi^{{
m Aut}(D)}|$ .

 $\Leftarrow$ : Denote  $\ell$  the isomorphism between  $\psi$  and  $\psi \circ \sigma$ . Define  $f: G \to G$  by  $vg \mapsto \ell(v)\sigma(g)$ . Then  $f \in \operatorname{Aut}(G)$  is such that  $f(x_1) = x_2$  and  $f(y_1) = y_2$ . Statement i is proved.

Note that  $\operatorname{Aut}(D)$  acts transitively on the set  $\mathcal{RM}_G$ . There is  $|\mathcal{RM}_G| = |\operatorname{Aut}(D)|/|\operatorname{Aut}(D)_\psi| = |\psi^{\operatorname{Aut}(D)}|$ , which completes the proof.

We claim that the number of G-rotary augmented PX maps equals  $|\psi^{{
m Aut}(D)}|$ .

r	$Irr(D_{2r})$	G	Γ	the count
r is odd	$\gamma_{1,1}$	$\mathbb{Z}_p \times D_{2r}$	$C^*(p, r, 0)$	1
	$\gamma_{-1,-1}$	$D_{2pr}$	$C^*(p,r,0,-1)$	1
	$\psi$ ( $d\geqslant$ 2)	$\mathbb{Z}_p^d \rtimes_{\psi} D_{2r}$	C(p,r,d-1)	$ \psi^{\mathrm{Aut}(D_{2r})} $
r is even	$\gamma_{1,1}$	$\mathbb{Z}_p \times D_{2r}$	$C^*(p, r, 0)$	1
	$\gamma_{1,-1},\gamma_{-1,1}$	$\mathbb{Z}_p \rtimes_{\gamma_{1,-1}} D_{2r}$	$C^*(p, r, 0)$	1
	$\gamma_{-1,-1}$	$D_{2pr}$	$C^*(p,r,0,-1)$	1
	$\psi$ ( $d\geqslant$ 2)	$\mathbb{Z}_p^d \rtimes_{\psi} D_{2r}$	C(p,r,d-1)	$ \psi^{\operatorname{Aut}(D_{2r})} $

Table 1: Irreducible G-rotary PX maps

The number of irreducible PX maps of length r

# [2, Theorem 5.7]

The number of irreducible rotary augmented PX map of length r is

- $\bigcirc$  |Irr( $D_{2r}$ )| if r is odd;
- $|\operatorname{Irr}(D_{2r})| 1 \text{ if } r \text{ is even.}$

# The number of *G*-rotary PX maps

 $\{G ext{-rotary augmented PX maps}\}\$ is in bijection with  $\{ ext{the }\operatorname{Aut}(G) ext{-orbits on rotary pairs }(\rho, au)\$ of G where  $|\rho|=2p\}$ 

# The number of G-rotary PX maps

{G-rotary augmented PX maps} is in bijection with {the  $\operatorname{Aut}(G)$ -orbits on rotary pairs  $(\rho,\tau)$  of G where  $|\rho|=2p$ }  $G=V:_{\psi}D=\mathbb{Z}_{p}^{d}:_{\psi}\mathrm{D}_{2r},\ d\geqslant 2,\ \psi$  is irreducible.

# The number of G-rotary PX maps

 $\{G ext{-rotary augmented PX maps}\}\$ is in bijection with  $\{ ext{the }\operatorname{Aut}(G) ext{-orbits on rotary pairs }(\rho,\tau)\$ of G where  $|\rho|=2p\}$   $G=V:_{\psi}D=\mathbb{Z}_p^d:_{\psi}\mathrm{D}_{2r},\ d\geqslant 2,\ \psi$  is irreducible.

## [2, Lemma 5.4]

The group G has

$$p^d(p^{d/2}-1)r\varphi(r)$$

rotary pairs  $(\rho, \tau)$  such that  $|\rho| = 2p$ .

#### Proof.

The element  $\rho$  equals vc for some  $c \in D$  and v not in the -1-eigenspace of c.

 $\{G ext{-rotary augmented PX maps}\}\$ is in bijection with  $\{ ext{the }\operatorname{Aut}(G) ext{-orbits on rotary pairs }(\rho,\tau) \text{ of } G \text{ where } |\rho|=2p\}$   $G=V:_{\psi}D=\mathbb{Z}_p^d:_{\psi}\mathrm{D}_{2r},\ d\geqslant 2,\ \psi \text{ is irreducible.}$ 

## [2, Lemma 5.4]

The group G has

$$p^d(p^{d/2}-1)r\varphi(r)$$

rotary pairs  $(\rho, \tau)$  such that  $|\rho| = 2p$ .

#### Proof.

The element  $\rho$  equals vc for some  $c \in D$  and v not in the -1-eigenspace of c. Since the -1-eigenspace of a none centering involution in D has dimension d/2, there are  $(p^d-p^{d/2})r$  choices of  $\rho$ .

 $\{G ext{-rotary augmented PX maps}\}\$ is in bijection with  $\{ ext{the }\operatorname{Aut}(G) ext{-orbits on rotary pairs }(\rho,\tau)\$ of G where  $|\rho|=2p\}$   $G=V:_{\psi}D=\mathbb{Z}_p^d:_{\psi}\mathrm{D}_{2r},\ d\geqslant 2,\ \psi$  is irreducible.

# [2, Lemma 5.4]

The group G has

$$p^d(p^{d/2}-1)r\varphi(r)$$

rotary pairs  $(\rho, \tau)$  such that  $|\rho| = 2p$ .

#### Proof.

The element  $\rho$  equals vc for some  $c\in D$  and v not in the -1-eigenspace of c. Since the -1-eigenspace of a none centering involution in D has dimension d/2, there are  $(p^d-p^{d/2})r$  choices of  $\rho$ .

The involution  $\tau$  equals wz for some  $z \in D$  and w in the -1-eigenspace of z.

{G-rotary augmented PX maps} is in bijection with {the  $\operatorname{Aut}(G)$ -orbits on rotary pairs  $(\rho, \tau)$  of G where  $|\rho| = 2p$ }  $G = V:_{\psi}D = \mathbb{Z}_p^d:_{\psi}\operatorname{D}_{2r}, \ d \geqslant 2, \ \psi$  is irreducible.

## [2, Lemma 5.4]

The group G has

$$p^d(p^{d/2}-1)r\varphi(r)$$

rotary pairs  $(\rho, \tau)$  such that  $|\rho| = 2p$ .

#### Proof.

The element  $\rho$  equals vc for some  $c\in D$  and v not in the -1-eigenspace of c. Since the -1-eigenspace of a none centering involution in D has dimension d/2, there are  $(p^d-p^{d/2})r$  choices of  $\rho$ .

The involution  $\tau$  equals wz for some  $z \in D$  and w in the -1-eigenspace of z. So there are  $p^{d/2}\varphi(r)$  choices of  $\tau$ .

$$G = V:_{\psi} D = \mathbb{Z}_p^d:_{\psi} D_{2r}, \ d \geqslant 2, \ \psi \ \text{is irreducible}.$$

$$G = V:_{\psi}D = \mathbb{Z}_p^d:_{\psi}D_{2r}, \ d \geqslant 2, \ \psi$$
 is irreducible.

$$A := \operatorname{Aut}(G)$$

$$A = \operatorname{Inn}_G(V) : \operatorname{N}_A(D).$$

$$G = V:_{\psi}D = \mathbb{Z}_p^d:_{\psi}D_{2r}, d \geqslant 2, \psi$$
 is irreducible.

$$A := \operatorname{Aut}(G)$$

$$A = \operatorname{Inn}_{G}(V) : \operatorname{N}_{A}(D).$$

The number of of G-rotary augmented PX maps equals

$$(p^{d/2}-1)r\varphi(r)/|\mathrm{N}_A(D)|.$$

$$G = V_{\cdot \psi} D = \mathbb{Z}_p^d_{\cdot \psi} D_{2r}, \ d \geqslant 2, \ \psi \text{ is irreducible.}$$

# $A := \operatorname{Aut}(G)$

$$A = \operatorname{Inn}_{G}(V) : \operatorname{N}_{A}(D).$$

The number of of G-rotary augmented PX maps equals

$$(p^{d/2}-1)r\varphi(r)/|\mathrm{N}_A(D)|.$$

# [2, Lemma 5.2]

Define a homomorphism  $\kappa: \mathrm{N}_A(D) \to \mathrm{Aut}(D)$  by  $f \mapsto f|_D$ . Then  $\ker(\kappa) \cong \mathrm{End}_D(V)^{\times}$  and

$$\operatorname{im}(\kappa) = \operatorname{Aut}(D)_{\psi},$$

where  $\operatorname{Aut}(D)_{\psi} = \{ \sigma \in \operatorname{Aut}(D) \mid \psi \circ \sigma \cong \psi \}.$ 

$$G = V:_{\psi}D = \mathbb{Z}_p^d:_{\psi}D_{2r}, \ d \geqslant 2, \ \psi \text{ is irreducible.}$$

## A := Aut(G)

$$A = \operatorname{Inn}_{G}(V) : \operatorname{N}_{A}(D).$$

The number of of G-rotary augmented PX maps equals

$$(p^{d/2}-1)r\varphi(r)/|\mathrm{N}_A(D)|.$$

## [2, Lemma 5.2]

Define a homomorphism  $\kappa: \mathrm{N}_A(D) \to \mathrm{Aut}(D)$  by  $f \mapsto f|_D$ . Then  $\ker(\kappa) \cong \mathrm{End}_D(V)^{\times}$  and

$$\operatorname{im}(\kappa) = \operatorname{Aut}(D)_{\psi},$$

where  $\operatorname{Aut}(D)_{\psi} = \{ \sigma \in \operatorname{Aut}(D) \mid \psi \circ \sigma \cong \psi \}.$ 

## Sketch of the proof

$$\ker(\kappa) = \{ \sigma \in N \mid \sigma \mid_D = \mathrm{id}_D \}.$$

$$G = V:_{\psi}D = \mathbb{Z}_p^d:_{\psi}D_{2r}, \ d \geqslant 2, \ \psi \text{ is irreducible.}$$

## A := Aut(G)

$$A = \operatorname{Inn}_{G}(V) : \operatorname{N}_{A}(D).$$

The number of of *G*-rotary augmented PX maps equals

$$(p^{d/2}-1)r\varphi(r)/|\mathrm{N}_A(D)|.$$

## [2, Lemma 5.2]

Define a homomorphism  $\kappa: \mathrm{N}_A(D) \to \mathrm{Aut}(D)$  by  $f \mapsto f|_D$ . Then  $\ker(\kappa) \cong \mathrm{End}_D(V)^{\times}$  and

$$\operatorname{im}(\kappa) = \operatorname{Aut}(D)_{\psi},$$

where  $\operatorname{Aut}(D)_{\psi} = \{ \sigma \in \operatorname{Aut}(D) \mid \psi \circ \sigma \cong \psi \}.$ 

## Sketch of the proof

 $\ker(\kappa) = \{ \sigma \in N \mid \sigma \mid_D = \mathrm{id}_D \}.$  For each  $\ell \in \mathrm{End}_D(V)^{\times}$ , define  $\sigma_{\ell} : G \to G$  by  $vb \mapsto \ell(v)b$ .

$$G = V:_{\psi}D = \mathbb{Z}_p^d:_{\psi}D_{2r}, \ d \geqslant 2, \ \psi \text{ is irreducible.}$$

## A := Aut(G)

$$A = \operatorname{Inn}_{G}(V) : \operatorname{N}_{A}(D).$$

The number of of *G*-rotary augmented PX maps equals

$$(p^{d/2}-1)r\varphi(r)/|\mathrm{N}_A(D)|.$$

### [2, Lemma 5.2]

Define a homomorphism  $\kappa: \mathrm{N}_A(D) \to \mathrm{Aut}(D)$  by  $f \mapsto f|_D$ . Then  $\ker(\kappa) \cong \mathrm{End}_D(V)^{\times}$  and

$$\operatorname{im}(\kappa) = \operatorname{Aut}(D)_{\psi},$$

where  $\operatorname{Aut}(D)_{\psi} = \{ \sigma \in \operatorname{Aut}(D) \mid \psi \circ \sigma \cong \psi \}.$ 

## Sketch of the proof

 $\ker(\kappa) = \{ \sigma \in N \mid \sigma \mid_D = \mathrm{id}_D \}. \text{For each } \ell \in \mathrm{End}_D(V)^\times, \text{ define } \sigma_\ell : G \to G \text{ by } vb \mapsto \ell(v)b. \text{Then } \ker(\kappa) \supseteq \{ \sigma_\ell \mid \ell \in \mathrm{End}_D(V)^\times \}.$ 

$$G = V:_{\psi}D = \mathbb{Z}_p^d:_{\psi}D_{2r}, \ d \geqslant 2, \ \psi \text{ is irreducible.}$$

## A := Aut(G)

$$A = \operatorname{Inn}_{G}(V) : \operatorname{N}_{A}(D).$$

The number of of *G*-rotary augmented PX maps equals

$$(p^{d/2}-1)r\varphi(r)/|\mathrm{N}_A(D)|.$$

### [2, Lemma 5.2]

Define a homomorphism  $\kappa: \mathrm{N}_A(D) \to \mathrm{Aut}(D)$  by  $f \mapsto f|_D$ . Then  $\ker(\kappa) \cong \mathrm{End}_D(V)^{\times}$  and

$$\operatorname{im}(\kappa) = \operatorname{Aut}(D)_{\psi},$$

where  $\operatorname{Aut}(D)_{\psi} = \{ \sigma \in \operatorname{Aut}(D) \mid \psi \circ \sigma \cong \psi \}.$ 

## Sketch of the proof

 $\ker(\kappa) = \{\sigma \in N \mid \sigma \mid_D = \mathrm{id}_D\}. \text{For each } \ell \in \mathrm{End}_D(V)^\times, \text{ define } \sigma_\ell : G \to G \text{ by } vb \mapsto \ell(v)b. \text{Then } \ker(\kappa) \supseteq \{\sigma_\ell \mid \ell \in \mathrm{End}_D(V)^\times\}. \text{One the other hand, the function } N \to \mathrm{End}_D(V)^\times, \ \sigma \mapsto \sigma \mid_V, \text{ is injective since } \sigma \mid_D = \mathrm{id}_D.$ 

## Sketch of the proof

For each  $\delta \in \operatorname{Aut}(D)_{\psi}$ , there exists  $\sigma \in \operatorname{Aut}(G)$  such that  $\sigma \mid_{D} = \delta$ .

## Sketch of the proof

For each  $\delta \in \operatorname{Aut}(D)_{\psi}$ , there exists  $\sigma \in \operatorname{Aut}(G)$  such that  $\sigma \mid_{D} = \delta$ . Fix such an automorphism  $\delta$  of D, then there exists an  $\mathbb{F}_{p}D$ -module isomorphism  $\delta_{V}$  such that  $\delta_{V}(\psi(b)(v)) = \psi \circ \delta(b)\delta_{V}(v)$  for each  $b \in D$  and  $v \in V$ . Define

$$\sigma: G \to G, \ vb \mapsto \delta_V(v)\delta(b),$$

which is an automorphism of G.

#### Sketch of the proof

For each  $\delta \in \operatorname{Aut}(D)_{\psi}$ , there exists  $\sigma \in \operatorname{Aut}(G)$  such that  $\sigma \mid_{D} = \delta$ . Fix such an automorphism  $\delta$  of D, then there exists an  $\mathbb{F}_p D$ -module isomorphism  $\delta_V$  such that  $\delta_V(\psi(b)(v)) = \psi \circ \delta(b)\delta_V(v)$  for each  $b \in D$  and  $v \in V$ . Define

$$\sigma: G \to G, \ vb \mapsto \delta_V(v)\delta(b),$$

which is an automorphism of G. Thus  $\operatorname{im}(\kappa) = \operatorname{Aut}(D)_{\psi}$ .

#### Sketch of the proof

For each  $\delta \in \operatorname{Aut}(D)_{\psi}$ , there exists  $\sigma \in \operatorname{Aut}(G)$  such that  $\sigma \mid_{D} = \delta$ . Fix such an automorphism  $\delta$  of D, then there exists an  $\mathbb{F}_p D$ -module isomorphism  $\delta_V$  such that  $\delta_V(\psi(b)(v)) = \psi \circ \delta(b)\delta_V(v)$  for each  $b \in D$  and  $v \in V$ . Define

$$\sigma: G \to G, \ vb \mapsto \delta_V(v)\delta(b),$$

which is an automorphism of G. Thus  $\operatorname{im}(\kappa) = \operatorname{Aut}(D)_{\psi}$ .

#### Sketch of the proof

For each  $\delta \in \operatorname{Aut}(D)_{\psi}$ , there exists  $\sigma \in \operatorname{Aut}(G)$  such that  $\sigma \mid_{D} = \delta$ . Fix such an automorphism  $\delta$  of D, then there exists an  $\mathbb{F}_p D$ -module isomorphism  $\delta_V$  such that  $\delta_V(\psi(b)(v)) = \psi \circ \delta(b)\delta_V(v)$  for each  $b \in D$  and  $v \in V$ . Define

$$\sigma: G \to G, \ vb \mapsto \delta_V(v)\delta(b),$$

which is an automorphism of G. Thus  $\operatorname{im}(\kappa) = \operatorname{Aut}(D)_{\psi}$ .

Since  $\operatorname{End}_{\mathcal{D}}(V) \cong \mathbb{F}_p^{d/2}$ , we have that the number of *G*-rotary augmented PX maps equals  $\varphi(r)/|\operatorname{N}_A(D)| = |\operatorname{Aut}(D)|/|\operatorname{Aut}(D)_{\psi} = |\psi^{\operatorname{Aut}(D)}|$ .

For each  $\delta \in \operatorname{Aut}(D)_{\psi}$ , there exists  $\sigma \in \operatorname{Aut}(G)$  such that  $\sigma \mid_{D} = \delta$ . Fix such an automorphism  $\delta$  of D, then there exists an  $\mathbb{F}_p D$ -module isomorphism  $\delta_V$  such that  $\delta_V(\psi(b)(v)) = \psi \circ \delta(b)\delta_V(v)$  for each  $b \in D$  and  $v \in V$ . Define

$$\sigma: G \to G, \ vb \mapsto \delta_V(v)\delta(b),$$

which is an automorphism of G. Thus  $\operatorname{im}(\kappa) = \operatorname{Aut}(D)_{\psi}$ .

Since  $\operatorname{End}_{\mathcal{D}}(V) \cong \mathbb{F}_p^{d/2}$ , we have that the number of *G*-rotary augmented PX maps equals  $\varphi(r)/|\operatorname{N}_A(D)| = |\operatorname{Aut}(D)|/|\operatorname{Aut}(D)_{\psi} = |\psi^{\operatorname{Aut}(D)}|$ .

## [2, Lemma 5.2]

For  $i \in \{1,2\}$ , let  $G_i = \mathbb{Z}_p^{d_i} :_{\psi_i} D_i$ , where  $p \nmid |D_1||D_2|$ . Then the following are equivalent:

- $\bullet$  there exists an isomorphism  $\sigma: D_1 \to D_2$  such that  $\psi_2 \circ \sigma \cong \psi_1$ .

Moreover, if (ii) holds, there exists an isomorphism  $f:G_1\to G_2$  such that  $f(D_1)=D_2$  and  $f|_{D_1}=\sigma$ .

Let  $\mathcal M$  be a rotary augmented PX map of length r. Then  $\mathcal M$  is isomorphic to a direct product of irreducible rotary augmented PX maps of length r. Conversely, every direct product of irreducible rotary augmented PX maps of length r is a rotary augmented PX map whose underlying graph is of length r.

Let  $\mathcal M$  be a rotary augmented PX map of length r. Then  $\mathcal M$  is isomorphic to a direct product of irreducible rotary augmented PX maps of length r. Conversely, every direct product of irreducible rotary augmented PX maps of length r is a rotary augmented PX map whose underlying graph is of length r.

By definition, for every rotary map  $\mathcal{M}$ , we have  $\mathcal{M} \times \mathcal{M} \cong \mathcal{M}$ .

Let  $\mathcal M$  be a rotary augmented PX map of length r. Then  $\mathcal M$  is isomorphic to a direct product of irreducible rotary augmented PX maps of length r. Conversely, every direct product of irreducible rotary augmented PX maps of length r is a rotary augmented PX map whose underlying graph is of length r.

By definition, for every rotary map  $\mathcal{M}$ , we have  $\mathcal{M} \times \mathcal{M} \cong \mathcal{M}$ . Therefore, if there exists a decomposition  $\mathcal{M} \cong \mathcal{N}_1 \times \cdots \times \mathcal{N}_n$  for an arc-regular map  $\mathcal{M}$ , then we can pick  $i_1, i_2, \ldots, i_k$  such that  $\mathcal{M} \cong \mathcal{N}_{i_1} \times \cdots \times \mathcal{N}_{i_k}$  and  $\mathcal{N}_{i_{j_1}} \not\cong \mathcal{N}_{i_{j_2}}$  if  $j_1 \neq j_2$ .

Let  $\mathcal M$  be a rotary augmented PX map of length r. Then  $\mathcal M$  is isomorphic to a direct product of irreducible rotary augmented PX maps of length r. Conversely, every direct product of irreducible rotary augmented PX maps of length r is a rotary augmented PX map whose underlying graph is of length r.

By definition, for every rotary map  $\mathcal{M}$ , we have  $\mathcal{M} \times \mathcal{M} \cong \mathcal{M}$ . Therefore, if there exists a decomposition  $\mathcal{M} \cong \mathcal{N}_1 \times \cdots \times \mathcal{N}_n$  for an arc-regular map  $\mathcal{M}$ , then we can pick  $i_1, i_2, \ldots, i_k$  such that  $\mathcal{M} \cong \mathcal{N}_{i_1} \times \cdots \times \mathcal{N}_{i_k}$  and  $\mathcal{N}_{i_{j_1}} \not\cong \mathcal{N}_{i_{j_2}}$  if  $j_1 \neq j_2$ .

## [2, Lemma 6.4]

Let  $V_1=\mathbb{Z}_p^{d_1}$  and let  $V_2=\mathbb{Z}_p^{d_2}$ . Let  $G=(V_1\times V_2):_{(\psi_1,\psi_2)}\mathsf{D}_{2r}$  where  $\psi_i:\mathsf{D}_{2r}\to\mathsf{GL}(V_i)$  is irreducible for every  $i\in\{1,2\}$ . If G has a rotary pair  $(\rho,\tau)$  such that  $|\rho|=2p$  and  $|\tau|=2$ , then  $\psi_1\not\cong\psi_2$ .

Suppose the contradiction. There are  $v \in V_1 \times V_2$ ,  $\tau_1 \in D_r$  such that  $\rho = v\tau_1$ .

Suppose the contradiction. There are  $v \in V_1 \times V_2$ ,  $\tau_1 \in D_r$  such that  $\rho = v\tau_1$ . Let  $v' = \rho^2$ , and  $\tau' = \rho^p$  which is an involution. Let  $D = \langle \tau, \tau_1 \rangle$ .

Suppose the contradiction. There are  $v \in V_1 \times V_2$ ,  $\tau_1 \in D_r$  such that  $\rho = v\tau_1$ . Let  $v' = \rho^2$ , and  $\tau' = \rho^p$  which is an involution. Let  $D = \langle \tau, \tau_1 \rangle$ . Then  $G = (V_1 \times V_2)D$  implying that  $D \cong D_{2pr}$  or  $D_{2r}$ .

Suppose the contradiction. There are  $v \in V_1 \times V_2$ ,  $\tau_1 \in \mathrm{D}_r$  such that  $\rho = v\tau_1$ . Let  $v' = \rho^2$ , and  $\tau' = \rho^p$  which is an involution. Let  $D = \langle \tau, \tau_1 \rangle$ . Then  $G = (V_1 \times V_2)D$  implying that  $D \cong D_{2pr}$  or  $\mathrm{D}_{2r}$ . If  $D \cong \mathrm{D}_{2pr}$ , then  $D \cap (V_1 \times V_2) \cong \mathbb{Z}_p$  is a  $D_{2r}$ -submodule. Thus  $V_1 \cong V_2 \cong \mathbb{Z}_p$ . Contradiction comes.

Suppose the contradiction. There are  $v \in V_1 \times V_2$ ,  $\tau_1 \in \mathrm{D}_r$  such that  $\rho = v\tau_1$ . Let  $v' = \rho^2$ , and  $\tau' = \rho^p$  which is an involution. Let  $D = \langle \tau, \tau_1 \rangle$ . Then  $G = (V_1 \times V_2)D$  implying that  $D \cong D_{2pr}$  or  $\mathrm{D}_{2r}$ . If  $D \cong \mathrm{D}_{2pr}$ , then  $D \cap (V_1 \times V_2) \cong \mathbb{Z}_p$  is a  $D_{2r}$ -submodule. Thus  $V_1 \cong V_2 \cong \mathbb{Z}_p$ . Contradiction comes. We therefore have that  $D \cong \mathrm{D}_{2r}$ . So

$$G=(V_1\times V_2):_{(\eta_1,\eta_2)}D.$$

Suppose the contradiction. There are  $v \in V_1 \times V_2$ ,  $\tau_1 \in \mathrm{D}_r$  such that  $\rho = v\tau_1$ . Let  $v' = \rho^2$ , and  $\tau' = \rho^p$  which is an involution. Let  $D = \langle \tau, \tau_1 \rangle$ . Then  $G = (V_1 \times V_2)D$  implying that  $D \cong D_{2pr}$  or  $\mathrm{D}_{2r}$ . If  $D \cong \mathrm{D}_{2pr}$ , then  $D \cap (V_1 \times V_2) \cong \mathbb{Z}_p$  is a  $D_{2r}$ -submodule. Thus  $V_1 \cong V_2 \cong \mathbb{Z}_p$ . Contradiction comes. We therefore have that  $D \cong \mathrm{D}_{2r}$ . So

$$G=(V_1\times V_2)_{(\eta_1,\eta_2)}D.$$

By Corollary19, there exists  $\sigma \in \operatorname{Aut}(D_{2r})$  such that  $\eta_1 = \psi_1 \circ \sigma$  and  $\eta_2 \cong \psi_2 \circ \sigma$ .

Suppose the contradiction. There are  $v \in V_1 \times V_2$ ,  $\tau_1 \in D_r$  such that  $\rho = v\tau_1$ . Let  $v' = \rho^2$ , and  $\tau' = \rho^p$  which is an involution. Let  $D = \langle \tau, \tau_1 \rangle$ . Then  $G = (V_1 \times V_2)D$  implying that  $D \cong D_{2pr}$  or  $D_{2r}$ . If  $D \cong D_{2pr}$ , then  $D \cap (V_1 \times V_2) \cong \mathbb{Z}_p$  is a  $D_{2r}$ -submodule. Thus  $V_1 \cong V_2 \cong \mathbb{Z}_p$ . Contradiction comes. We therefore have that  $D \cong D_{2r}$ . So

$$G=(V_1\times V_2):_{(\eta_1,\eta_2)}D.$$

By Corollary19, there exists  $\sigma \in \operatorname{Aut}(\mathbb{D}_{2r})$  such that  $\eta_1 = \psi_1 \circ \sigma$  and  $\eta_2 \cong \psi_2 \circ \sigma$ . So  $\eta_1 \cong \eta_2$ , denote by f this D-module isomorphism. Let  $v' = (x,y) \in V_1 \times V_2$ . Then  $x \in \operatorname{C}_{V_1}(\tau')$  and  $y \in \operatorname{C}_{V_2}(\tau')$ .

Suppose the contradiction. There are  $v \in V_1 \times V_2$ ,  $\tau_1 \in D_r$  such that  $\rho = v\tau_1$ . Let  $v' = \rho^2$ , and  $\tau' = \rho^p$  which is an involution. Let  $D = \langle \tau, \tau_1 \rangle$ . Then  $G = (V_1 \times V_2)D$  implying that  $D \cong D_{2pr}$  or  $D_{2r}$ . If  $D \cong D_{2pr}$ , then  $D \cap (V_1 \times V_2) \cong \mathbb{Z}_p$  is a  $D_{2r}$ -submodule. Thus  $V_1 \cong V_2 \cong \mathbb{Z}_p$ . Contradiction comes. We therefore have that  $D \cong D_{2r}$ . So

$$G=(V_1\times V_2)_{(\eta_1,\eta_2)}D.$$

By Corollary19, there exists  $\sigma \in \operatorname{Aut}(D_{2r})$  such that  $\eta_1 = \psi_1 \circ \sigma$  and  $\eta_2 \cong \psi_2 \circ \sigma$ . So  $\eta_1 \cong \eta_2$ , denote by f this D-module isomorphism. Let  $v' = (x,y) \in V_1 \times V_2$ . Then  $x \in \operatorname{C}_{V_1}(\tau')$  and  $y \in \operatorname{C}_{V_2}(\tau')$ . By Lemma[2, Lemma 2.10], there exists  $\delta' \in \operatorname{End}_D(V_1)$  and  $y' \in V_1$  such that

$$\delta'(x) = y' = f^{-1}(y).$$

Suppose the contradiction. There are  $v \in V_1 \times V_2$ ,  $\tau_1 \in D_r$  such that  $\rho = v\tau_1$ . Let  $v' = \rho^2$ , and  $\tau' = \rho^p$  which is an involution. Let  $D = \langle \tau, \tau_1 \rangle$ . Then  $G = (V_1 \times V_2)D$  implying that  $D \cong D_{2pr}$  or  $D_{2r}$ . If  $D \cong D_{2pr}$ , then  $D \cap (V_1 \times V_2) \cong \mathbb{Z}_p$  is a  $D_{2r}$ -submodule. Thus  $V_1 \cong V_2 \cong \mathbb{Z}_p$ . Contradiction comes. We therefore have that  $D \cong D_{2r}$ . So

$$G = (V_1 \times V_2):_{(\eta_1, \eta_2)} D.$$

By Corollary19, there exists  $\sigma \in \operatorname{Aut}(D_{2r})$  such that  $\eta_1 = \psi_1 \circ \sigma$  and  $\eta_2 \cong \psi_2 \circ \sigma$ . So  $\eta_1 \cong \eta_2$ , denote by f this D-module isomorphism. Let  $v' = (x,y) \in V_1 \times V_2$ . Then  $x \in \operatorname{C}_{V_1}(\tau')$  and  $y \in \operatorname{C}_{V_2}(\tau')$ . By Lemma[2, Lemma 2.10], there exists  $\delta' \in \operatorname{End}_D(V_1)$  and  $y' \in V_1$  such that

$$\delta'(x) = y' = f^{-1}(y).$$

Define  $\delta' = \delta \circ f$ , which is a *D*-module isomorphism such that  $\delta(x) = y$ . Therefore,  $v' \in \{(w, \zeta(w)) \mid w \in V_1\}$ .

Suppose the contradiction. There are  $v \in V_1 \times V_2$ ,  $\tau_1 \in D_r$  such that  $\rho = v\tau_1$ . Let  $v' = \rho^2$ , and  $\tau' = \rho^p$  which is an involution. Let  $D = \langle \tau, \tau_1 \rangle$ . Then  $G = (V_1 \times V_2)D$  implying that  $D \cong D_{2pr}$  or  $D_{2r}$ . If  $D \cong D_{2pr}$ , then  $D \cap (V_1 \times V_2) \cong \mathbb{Z}_p$  is a  $D_{2r}$ -submodule. Thus  $V_1 \cong V_2 \cong \mathbb{Z}_p$ . Contradiction comes. We therefore have that  $D \cong D_{2r}$ . So

$$G = (V_1 \times V_2):_{(\eta_1, \eta_2)} D.$$

By Corollary19, there exists  $\sigma \in \operatorname{Aut}(D_{2r})$  such that  $\eta_1 = \psi_1 \circ \sigma$  and  $\eta_2 \cong \psi_2 \circ \sigma$ . So  $\eta_1 \cong \eta_2$ , denote by f this D-module isomorphism. Let  $v' = (x,y) \in V_1 \times V_2$ . Then  $x \in \operatorname{C}_{V_1}(\tau')$  and  $y \in \operatorname{C}_{V_2}(\tau')$ . By Lemma[2, Lemma 2.10], there exists  $\delta' \in \operatorname{End}_D(V_1)$  and  $y' \in V_1$  such that

$$\delta'(x) = y' = f^{-1}(y).$$

Define  $\delta' = \delta \circ f$ , which is a D-module isomorphism such that  $\delta(x) = y$ . Therefore,  $v' \in \{(w, \zeta(w)) \mid w \in V_1\}$ . Note that  $\{(w, \zeta(w)) \mid w \in V_1\} \cong \mathbb{Z}_p^{d_1}$  is normal in G. We have that  $\langle (v')^D \rangle \leqslant \{(w, \zeta(w)) \mid w \in V_1\}$ .

Suppose the contradiction. There are  $v \in V_1 \times V_2$ ,  $\tau_1 \in D_r$  such that  $\rho = v\tau_1$ . Let  $v' = \rho^2$ , and  $\tau' = \rho^p$  which is an involution. Let  $D = \langle \tau, \tau_1 \rangle$ . Then  $G = (V_1 \times V_2)D$  implying that  $D \cong D_{2pr}$  or  $D_{2r}$ . If  $D \cong D_{2pr}$ , then  $D \cap (V_1 \times V_2) \cong \mathbb{Z}_p$  is a  $D_{2r}$ -submodule. Thus  $V_1 \cong V_2 \cong \mathbb{Z}_p$ . Contradiction comes. We therefore have that  $D \cong D_{2r}$ . So

$$G = (V_1 \times V_2):_{(\eta_1, \eta_2)} D.$$

By Corollary19, there exists  $\sigma \in \operatorname{Aut}(D_{2r})$  such that  $\eta_1 = \psi_1 \circ \sigma$  and  $\eta_2 \cong \psi_2 \circ \sigma$ . So  $\eta_1 \cong \eta_2$ , denote by f this D-module isomorphism. Let  $v' = (x,y) \in V_1 \times V_2$ . Then  $x \in \operatorname{C}_{V_1}(\tau')$  and  $y \in \operatorname{C}_{V_2}(\tau')$ . By Lemma[2, Lemma 2.10], there exists  $\delta' \in \operatorname{End}_D(V_1)$  and  $y' \in V_1$  such that

$$\delta'(x) = y' = f^{-1}(y).$$

Define  $\delta' = \delta \circ f$ , which is a D-module isomorphism such that  $\delta(x) = y$ . Therefore,  $v' \in \{(w, \zeta(w)) \mid w \in V_1\}$ . Note that  $\{(w, \zeta(w)) \mid w \in V_1\} \cong \mathbb{Z}_p^{d_1}$  is normal in G. We have that  $\langle (v')^D \rangle \leqslant \{(w, \zeta(w)) \mid w \in V_1\}$ . However, this is impossible because  $\{(w, \zeta(w)) \mid w \in V_1\} \neq V_1 \times V_2$ . Therefore, we have that  $\psi_1 \not\cong \psi_2$ .

#### Theorem

Let  $\mathcal{M}_1, \ldots, \mathcal{M}_m, \mathcal{N}_1, \ldots, \mathcal{N}_n$  be irreducible rotary augmented PX maps whose underlying graph is of length r. Let  $\mathcal{M}_i \ncong \mathcal{M}_j$  for  $i \neq j$  and let  $\mathcal{N}_i \ncong \mathcal{N}_j$  for  $i \neq j$ . Then  $\mathcal{M}_1 \times \cdots \times \mathcal{M}_m \cong \mathcal{N}_1 \times \cdots \times \mathcal{N}_n$  if and only if m = n and there is a permutation  $\sigma$  of  $\{1, \ldots, n\}$  such that  $\mathcal{M}_i \cong \mathcal{N}_{\sigma(i)}$  for all i.

- Map
- Rotary Praeger-Xu maps (PX maps)
  - Praeger-Xu graphs
  - The characterization of arc-regular automorphism groups of Praeger-Xu graphs
- Irreducible rotary PX maps
  - Construction of a class of irreducible rotary PX maps of length r
  - ullet The correspondence between irreducible rotary PX maps of length r and irreducible representations of  $D_{2r}$
  - ullet The count of  $V:_{\psi}\mathrm{D}_{2r}$ -rotary PX maps
- The decomposition of rotary PX maps

- [1] J. Chen, W. Fan, C. H. Li, and Y. Z. Zhu. Coverings of groups, regular dessins, and surfaces, 2024.
- [2] Z. Ding, Z. Guo, and L. Liu.Embedding a praeger-xu graph into a surface, 2025.
- [3] S. F. Du, G. Jones, J. H. Kwak, R. Nedela, and M. Škoviera. Regular embeddings of K<sub>n,n</sub> where n is a power of 2. I. Metacyclic case. *European Journal of Combinatorics*, 28(6):1595–1609, 2007.
- [4] S. F. Du, G. Jones, J. H. Kwak, R. Nedela, and M. Skoviera. Regular embeddings of K<sub>n,n</sub> where n is a power of 2. II: The non-metacyclic case. *European Journal of Combinatorics*, 31(7):1946–1956, 2010.
- [5] M. Giudici and G. Verret. Arc-transitive graphs of valency twice a prime admit a semiregular automorphism. Ars Mathematica Contemporanea, 18(1):179–186, 2020.
- [6] D. Gorenstein.
   Finite groups.
   Chelsea Publishing Co., New York, second edition, 1980.
- [7] Š. Gyürki, S. Pavlíková, and J. Širáň.
   Orientably-regular embeddings of complete multigraphs.

- J. Combin. Theory Ser. B, 171:71-95, 2025.
- [8] I. Martin Isaacs.
   *Finite Group Theory*.
   Number v. 92 in Graduate Studies in Mathematics. American Mathematical Society, Providence, R.I., 2008.
- [9] R. Jajcay, P. Potočnik, and S. Wilson. The praeger-xu graphs: Cycle structures, maps, and semitransitive orientations.

Acta Mathematica Universitatis Comenianae, 88(2):269–291, June 2019.

- [10] R. Jajcay, P. Potočnik, and S. Wilson. On the Cayleyness of Praeger-Xu graphs. Journal of Combinatorial Theory, Series B, 152:55–79, 2022.
- [11] L. D. James and G. A. Jones. Regular orientable imbeddings of complete graphs. Journal of Combinatorial Theory, Series B, 39(3):353–367, 1985.
- [12] G. A. Jones. Regular embeddings of complete bipartite graphs: Classification and enumeration. Proceedings of the London Mathematical Society. Third Series, 101(2):427–453, 2010.

- [13] G. A. Jones, R. Nedela, and M. Škoviera. Regular embeddings of K<sub>n,n</sub> where n is an odd prime power. European Journal of Combinatorics, 28(6):1863–1875, 2007.
- [14] S. K. Lando and A. K. Zvonkin. Graphs on Surfaces and Their Applications, volume 141 of Encyclopaedia of Mathematical Sciences. Springer Berlin Heidelberg, Berlin, Heidelberg, 2004.
- [15] C. H. Li, C. E. Praeger, and S. J. Song. Locally finite vertex-rotary maps and coset graphs with finite valency and finite edge multiplicity. *Journal of Combinatorial Theory. Series B*, 169:1–44, November 2024.
- [16] C. H. Li and Y. Z. Zhu. Covers and pseudocovers of symmetric graphs. Journal of Algebraic Combinatorics, 59(4):1021–1036, 2024.
- [17] P. Potočnik, P. Spiga, and G. Verret. Bounding the order of the vertex-stabiliser in 3-valent vertex-transitive and 4-valent arc-transitive graphs. Journal of Combinatorial Theory, Series B, 111:148–180, 2015.
- [18] C. E. Praeger.Highly Arc Transitive Digraphs.

- European Journal of Combinatorics, 10(3):281–292, 1989.
- [19] C. E. Praeger and M. Y. Xu. A Characterization of a Class of Symmetric Graphs of Twice Prime Valency. European Journal of Combinatorics, 10(1):91–102, 1989.
- [20] J. H. Van Lint. Introduction to Coding Theory, volume 86 of Graduate Texts in Mathematics. Springer Berlin Heidelberg, Berlin, Heidelberg, 1999.
- [21] J. Širáň, T. W. Tucker, and M. E. Watkins. Realizing finite edge-transitive orientable maps. J. Graph Theory, 37(1):1–34, 2001.